



KIT FOR THE PREPARATION OF TECHNETIUM TC 99M SESTAMIBI INJECTION

tetrakis (2-methoxy isobutyl isocyanide) copper(1) tetrafluoroborate injection, powder, lyophilized, for solution
JUBILANT DRAXIMAGE (USA) INC.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Kit for the Preparation of Technetium Tc 99m Sestamibi Injection safely and effectively. See full prescribing information for Kit for the Preparation of Technetium Tc 99m Sestamibi Injection.

Kit for the Preparation of Technetium Tc 99m Sestamibi Injection (tetrakis(1-isocyanato-2-methoxy-2-methyl-propane)-copper(1) tetrafluoroborate) injection, powder, lyophilized, for solution for Intravenous use.

Initial U.S. Approval: December 1990

INDICATIONS AND USAGE

Technetium Tc 99m Sestamibi is a myocardial perfusion agent indicated for:

- detecting coronary artery disease by localizing myocardial ischemia (reversible defects) and infarction (non-reversible defects) (1)
- evaluating myocardial function and developing information for use in patient management decisions (1)

DOSAGE AND ADMINISTRATION

For Myocardial Imaging: The suggested dose range for I.V. administration of Technetium Tc 99m Sestamibi in a single dose to be employed in the average patient (70 kg) is 370 to 1110 MBq (10 to 30 mCi) (2).

For Breast Imaging: The recommended dose range for I.V. administration of Technetium Tc 99m Sestamibi is a single dose of 740 to 1110 MBq (20 to 30 mCi) (2).

DOSAGE FORMS AND STRENGTHS

Kit for the Preparation of Technetium Tc 99m Sestamibi Injection is supplied as a lyophilized sterile and non-pyrogenic mixture in a 10 mL vial in kits of five (5) (NDC # 45548-141-05), ten (10) (NDC # 45548-141-10) and thirty (30) (NDC # 45548-141-30) (3).

CONTRAINDICATIONS

None known (4)

WARNINGS AND PRECAUTIONS

- Pharmacologic induction of cardiovascular stress may be associated with serious adverse events such as myocardial infarction, arrhythmia, hypotension, bronchoconstriction and cerebrovascular events (5.1).
- Technetium Tc 99m Sestamibi has been rarely associated with acute severe allergic and anaphylactic events of angioedema and generalized urticaria. In some patients the allergic symptoms developed on the second injection during Technetium Tc 99m Sestamibi imaging (5.1).
- Caution should be exercised and emergency equipment should be available when administering Technetium Tc 99m Sestamibi (5.1).
- Before administering Technetium Tc 99m Sestamibi patients should be asked about the possibility of allergic reactions to either drug (5.1).
- The contents of the vial are intended only for use in the preparation of Technetium Tc 99m Sestamibi and are not to be administered directly to the patient without first undergoing the preparative procedure (5.2).

ADVERSE REACTIONS

The following adverse reactions have been reported in $\leq 0.5\%$ of patients: signs and symptoms consistent with seizure occurring shortly after administration of the agent; transient arthritis, angioedema, arrhythmia, dizziness, syncope, abdominal pain, vomiting, and severe hypersensitivity characterized by dyspnea, hypotension, bradycardia, asthenia, and vomiting within two hours after a second injection of Technetium Tc 99m Sestamibi. A few cases of flushing, edema, injection site inflammation, dry mouth, fever, pruritis, rash, urticaria and fatigue have also been attributed to administration of the agent (6).

To report SUSPECTED ADVERSE REACTIONS, contact Jubilant DraxImage Inc. at 1-888-633-5343 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Specific drug-drug interactions have not been studied (7).

USE IN SPECIFIC POPULATIONS

- In one study of 46 subjects who received Technetium Tc 99m Sestamibi administration, the radioactivity in both children and adolescents exhibited blood PK profiles similar to those previously reported in adults (8.4).
- **Lactation:** Interruption of breastfeeding after exposure to Technetium Tc 99m Sestamibi is not necessary, however, a lactating woman should be advised to consider restricting close contact with her breast fed infant to a maximum of 5 hours in the 24 hour period after Technetium Tc 99m Sestamibi administration in order to minimize radiation exposure (8.2).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 3/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- Image Acquisition
- Radiation Dosimetry
- Instructions For Preparation
- Determination of Radiochemical Purity in Technetium Tc99m Sestamibi

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- Warnings
- General Precautions

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Lactation
- Pediatric Use
- Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

- Physical Characteristics
- External Radiation

12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Myocardial Imaging: Kit for the preparation of Technetium Tc 99m Sestamibi Injection is a myocardial perfusion agent that is indicated for detecting coronary artery disease by localizing myocardial ischemia (reversible defects) and infarction (non-reversible defects), in evaluating myocardial function and developing information for use in patient management decisions. Technetium Tc 99m Sestamibi evaluation of myocardial ischemia can be accomplished with rest and cardiovascular stress techniques (e.g., exercise or pharmacologic stress in accordance with the pharmacologic stress agent's labeling).

It is usually not possible to determine the age of a myocardial infarction or to differentiate a recent myocardial infarction from ischemia.

Breast Imaging: Kit for the preparation of Technetium Tc 99m Sestamibi Injection is indicated for planar imaging as a second line diagnostic drug after mammography to assist in the evaluation of breast lesions in patients with an abnormal mammogram or a palpable breast mass.

Kit for the preparation of Technetium Tc 99m Sestamibi Injection is not indicated for breast cancer screening, to confirm the presence or absence of malignancy, and it is not an alternative to biopsy.

2 DOSAGE AND ADMINISTRATION

For Myocardial Imaging: The suggested dose range for I.V. administration of Technetium Tc 99m Sestamibi in a single dose to be employed in the average patient (70 kg) is 370 to 1110 MBq (10 to 30 mCi).

For Breast Imaging: The recommended dose range for I.V. administration of Technetium Tc 99m Sestamibi is a single dose of 740 to 1110 MBq (20 to 30 mCi).

2.1 Image Acquisition

Breast Imaging: It is recommended that images be obtained with a table overlay to separate breast tissue from the myocardium and liver, and to exclude potential activity that may be present in the opposite breast. For lateral images, position the patient prone with the ipsilateral arm comfortably above the head, shoulders flat against the table, head turned to the side and relaxed, with the breast imaged pendent through an overlay cutout. The breast should not be compressed on the overlay. For anterior images, position the patient supine with both arms behind the head. For either lateral or anterior images, shield the chest and abdominal organs, or remove them from the field of view.

For complete study, sets of images should be obtained five minutes after the injection, and in the following sequence:

Beginning five minutes after the injection of Technetium Tc 99m Sestamibi:

- ten-minute lateral image of breast with abnormality
- ten-minute lateral image of contralateral breast
- ten-minute anterior image of both breasts

2.2 Radiation Dosimetry

The radiation doses to organs and tissues of an average patient (70 kg) per 1110 MBq (30 mCi) of Technetium Tc 99m Sestamibi injected intravenously are shown in Table 1.

Table 1. Radiation Absorbed Doses from Tc 99m Sestamibi

Organ	Estimated Radiation Absorbed Dose							
	REST				STRESS			
	2.0 hour void		4.8 hour void		2.0 hour void		4.8 hour void	
	rads/ 30 mCi	mGy/ 1110 MBq	rads/ 30 mCi	mGy/ 1110 MBq	rads/ 30 mCi	mGy/ 1110 MBq	rads/ 30 mCi	mGy/ 1110 MBq
Breasts	0.2	2.0	0.2	1.9	0.2	2.0	0.2	1.8
Gallbladder Wall	2.0	20.0	2.0	20.0	2.8	28.9	2.8	27.8
Small Intestine	3.0	30.0	3.0	30.0	2.4	24.4	2.4	24.4
Upper Large Intestine Wall	5.4	55.5	5.4	55.5	4.5	44.4	4.5	44.4
Lower Large Intestine Wall	3.9	40.0	4.2	41.1	3.3	32.2	3.3	32.2
Stomach Wall	0.6	6.1	0.6	5.8	0.6	5.3	0.5	5.2
Heart Wall	0.5	5.1	0.5	4.9	0.5	5.6	0.5	5.3
Kidneys	2.0	20.0	2.0	20.0	1.7	16.7	1.7	16.7
Liver	0.6	5.8	0.6	5.7	0.4	4.2	0.4	4.1
Lungs	0.3	2.8	0.3	2.7	0.3	2.6	0.2	2.4
Bone Surfaces	0.7	6.8	0.7	6.4	0.6	6.2	0.6	6.0
Thyroid	0.7	7.0	0.7	2.4	0.3	2.7	0.2	2.4
Ovaries	1.5	15.5	1.6	15.5	1.2	12.2	1.3	13.3
Testes	0.3	3.4	0.4	3.9	0.3	3.1	0.3	3.4
Red Marrow	0.5	5.1	0.5	5.0	0.5	4.6	0.5	4.4
Urinary Bladder Wall	2.0	20.0	4.2	41.1	1.5	15.5	3.0	30.0
Total Body	0.5	4.8	0.5	4.8	0.4	4.2	0.4	4.2

Radiation dosimetry calculations performed by Radiation Internal Dose Information Center, Oak Ridge Institute for Science and Education, PO Box 117, Oak Ridge, TN 37831-0117.

2.3 Instructions For Preparation

Preparation of the Technetium Tc 99m Sestamibi from the Kit for the Preparation of Technetium Tc 99m Sestamibi is done by the following aseptic procedure:

General Procedure:

- Prior to adding the Sodium Pertechnetate Tc 99m Injection to the vial, inspect the vial carefully for the presence of damage, particularly cracks, and do not use the vial if found. Tear off radiation symbol tag and attach it to the neck of the vial.
- Waterproof gloves should be worn during the preparation procedure. Remove the plastic disc from the vial and swab the top of the vial closure with alcohol to sanitize the surface.

Boiling Water Bath Procedure:

- Place the vial in a suitable radiation shield with a fitted radiation cap.
- With a sterile shielded syringe, aseptically obtain additive-free, sterile, non-pyrogenic Sodium Pertechnetate Tc 99m Injection [925 to 5550 MBq, (25 to 150 mCi)] in approximately 1 to 3 mL.
- Aseptically add the Sodium Pertechnetate Tc 99m Injection to the vial in the lead shield. Without withdrawing the needle, remove an equal volume of headspace to maintain atmospheric pressure within the vial.
- Shake vigorously, about 5 to 10 quick upward-downward motions.
- Remove the vial from the lead shield and place upright in an appropriately shielded and contained boiling water bath, such that the vial is suspended above the bottom of the bath, and boil for 10 minutes. Timing for 10 minutes is begun as soon as the water begins to boil again. Do not allow the boiling water to come in contact with the aluminum crimp.
- Remove the vial from the water bath, place in the lead shield and allow to cool for fifteen (15) minutes.
- Using proper shielding, the vial contents should be visually inspected. Use only if the solution is clear and free of particulate matter and discoloration.
- Assay the reaction vial using a suitable radioactivity calibration system. Record the Technetium Tc 99m concentration, total volume, assay time and date, expiration time and lot number on the radioassay information label and affix the label to the shield.
- Store the reaction vial containing the Technetium Tc 99m Sestamibi at 15 °C to 25 °C (59 °F to 77 °F) until use; at such time the product should be aseptically withdrawn. Technetium Tc 99m Sestamibi should be used within six (6) hours of preparation. The vial contains no preservative.

Note: Adherence to the above product reconstitution instructions is recommended.

The potential for cracking and significant contamination exists whenever vials containing radioactive material are heated.

This product is not to be used with the Recon-o-stat (thermal cycler) due to the smaller vial size requirements of this heating device.

Product should be used within six (6) hours after preparation.


Final product with radiochemical purity of at least 90 % was used in the clinical trials that established safety and effectiveness. The radiochemical purity was determined by the following method.

2.4 Determination of Radiochemical Purity in Technetium Tc 99m Sestamibi

- Obtain a Baker-Flex Aluminum Oxide coated, plastic TLC plate, #1 B-F, pre-cut to 2.5 cm x 7.5 cm.
- Dry the plate or plates at 100°C for 1 hour and store in a desiccator. Remove pre-dried plate from the desiccator just prior to use.
- Apply 1 drop of ethanol* using a 1 mL syringe with a 22 to 26 gauge needle, 1.5 cm from the bottom of the plate. THE SPOT SHOULD NOT BE ALLOWED TO DRY.
- Add 2 drops of Technetium Tc 99m Sestamibi solution, side by side on top of the ethanol* spot. Return the plate to a desiccator and allow the sample spot to dry (typically 15 minutes).
- The TLC tank is prepared by pouring ethanol* to a depth of 3 to 4 mm. Cover the tank and let it equilibrate for -10 minutes.
- Develop the plate in the covered TLC tank in ethanol* for a distance of 5 cm from the point of application.
- Cut the TLC plate 4 cm from the bottom and measure the Tc 99m activity in each piece by appropriate radiation detector.
- Calculate the % Tc 99m Sestamibi as:

$$\% \text{ Tc 99m Sestamibi} = \frac{\text{mCi Top Piece}}{\text{mCi Both Pieces}} \times 100$$

Figure 1. TLC Plate Diagram



*The ethanol used in this procedure should be 95 % or greater. Absolute ethanol (99 %) should remain at $\geq 95\%$ ethanol content for one week after opening if stored tightly capped, in a cool dry place.

3 DOSAGE FORMS AND STRENGTHS

Kit for Preparation of Technetium Tc 99m Sestamibi Injection is supplied as a lyophilized sterile and non-pyrogenic mixture in a 10 mL vial.

4 CONTRAINDICATIONS

None known.

5 WARNINGS AND PRECAUTIONS

5.1 Warnings

In studying patients in whom cardiac disease is known or suspected, care should be taken to assure continuous monitoring and treatment in accordance with safe, accepted clinical procedure. Infrequently, death has occurred 4 to 24 hours after Tc 99m Sestamibi use and is usually associated with exercise stress testing [see *General Precautions (5.2)*].

Pharmacologic induction of cardiovascular stress may be associated with serious adverse events such as myocardial infarction, arrhythmia, hypotension, bronchoconstriction and cerebrovascular events. Caution should be used when pharmacologic stress is selected as an alternative to exercise; it should be used when indicated and in accordance with the pharmacologic stress agent's labeling.

Technetium Tc 99m Sestamibi has been rarely associated with acute severe allergic and anaphylactic events of angioedema and generalized urticaria. In some patients the allergic symptoms developed on the second injection during Tc 99m Sestamibi imaging. Patients who receive Technetium Tc 99m Sestamibi for either myocardial or breast imaging are receiving the same drug. Caution should be exercised and emergency equipment should be available when administering Technetium Tc 99m Sestamibi. Also, before administering Technetium Tc 99m Sestamibi Injection, patients should be asked about the possibility of allergic reactions to the drug.

5.2 General Precaution

The contents of the vial are intended only for use in the preparation of Technetium Tc 99m Sestamibi and are not to be administered directly to the patient without first undergoing the preparative procedure.

Radioactive drugs must be handled with care and appropriate safety measures should be used to minimize radiation exposure to clinical personnel. Also, care should be taken to minimize radiation exposure to the patients consistent with proper patient management.

Contents of the kit before preparation are not radioactive. However, after the Sodium Pertechnetate Tc 99m Injection is added, adequate shielding of the final preparation must be maintained. The components of the kit are sterile and nonpyrogenic. It is essential to follow directions carefully and to adhere to strict aseptic procedures during preparation.

Technetium Tc 99m labeling reactions depend on maintaining the stannous ion in the reduced state. Hence, Sodium Pertechnetate Tc 99m Injection containing oxidants should not be used.

Technetium Tc 99m Sestamibi should not be used more than six (6) hours after preparation.

Radiopharmaceuticals should be used only by physicians who are qualified by training and experience in the safe use and handling of radionuclides and whose experience and training have been approved by the appropriate government agency authorized to license the use of radionuclides.

Stress testing should be performed only under the supervision of a qualified physician and in a laboratory equipped with appropriate resuscitation and support apparatus.

The most frequent exercise stress test endpoints sufficient to stop the test reported during controlled studies (two-thirds were cardiac patients) were:

Fatigue	35 %
Dyspnea	17 %
Chest Pain	16 %
ST-depression	7 %
Arrhythmia	1 %

6 ADVERSE REACTIONS

Adverse events were evaluated in 3741 adults who were evaluated in clinical studies. Of these patients, 3068 (77 % men, 22 % women, and 0.7 % of the patients' genders were not recorded) were in cardiac clinical trials and 673 (100 % women) in breast imaging trials. Cases of angina, chest pain, and death have occurred [see *Warnings and Precautions (5)*]. Adverse events reported at a rate of 0.5 % or greater after receiving Technetium Tc 99m Sestamibi administration are shown in the following table:

Table 2. Selected Adverse Events Reported in > 0.5 % of Patients Who Received Technetium Tc 99m Sestamibi in Either Breast or Cardiac Clinical Studies*

Body System	Breast studies		Cardiac studies	
	Women n=673	Women n=685	Men n=2361	Total n=3045
Body as a Whole	21 (3.1 %)	6 (0.9 %)	17 (0.7 %)	23 (0.8 %)
Headache	11 (1.6 %)	2 (0.3 %)	4 (0.2 %)	6 (0.2 %)
Cardiovascular	9 (1.3 %)	24 (3.5 %)	75 (3.2 %)	99 (3.3 %)
Chest Pain/Angina	0 (0 %)	18 (2.6 %)	46 (1.9 %)	64 (2.1 %)
ST segment changes	0 (0 %)	11 (1.6 %)	29 (1.2 %)	40 (1.3 %)
Digestive System	8 (1.2 %)	4 (0.6 %)	9 (0.4 %)	13 (0.4 %)
Nausea	4 (0.6 %)	1 (0.1 %)	2 (0.1 %)	3 (0.1 %)
Special Senses	132 (19.6 %)	62 (9.1 %)	160 (6.8 %)	222 (7.3 %)
Taste Perversion	129 (19.2 %)	60 (8.8 %)	157 (6.6 %)	217 (7.1 %)
Parosmia	8 (1.2 %)	6 (0.9 %)	10 (0.4 %)	16 (0.5 %)

*Excludes the 22 patients whose gender was not recorded.

In the clinical studies for breast imaging, breast pain was reported in 12 (1.7 %) of the patients. In 11 of these patients the pain appears to be associated with biopsy/surgical procedures.

The following adverse reactions have been reported in $\leq 0.5\%$ of patients: signs and symptoms consistent with seizure occurring shortly after administration of the agent; transient arthritis; angioedema, arrhythmia, dizziness, syncope, abdominal pain, vomiting, and severe hypersensitivity characterized by dyspnea, hypotension, bradycardia, asthenia, and vomiting within two hours after a second injection of Technetium Tc 99m Sestamibi. A few cases of flushing, edema, injection site inflammation, dry mouth, fever, pruritis, rash, urticaria and fatigue have also been attributed to administration of the agent.

7 DRUG INTERACTIONS

Specific drug-drug interactions have not been studied.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited available data with Technetium Tc 99m Sestamibi use in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Animal reproduction and teratogenicity studies have not been conducted with Technetium Tc 99m Sestamibi. However, all radiopharmaceuticals have the potential to cause fetal harm depending on the fetal stage of development and the magnitude of the radiation dose. If considering Technetium

Tc 99m Sestamibi administration to a pregnant woman, inform the patient about the potential for adverse pregnancy outcomes based on the radiation dose from Technetium Tc 99m Sestamibi and the gestational timing of exposure.

All pregnancies have a background risk of birth defects, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 % to 4 % and 15 % to 20 %, respectively.

8.2 Lactation

Risk Summary

Limited data in the scientific literature on the presence of Technetium Tc 99m Sestamibi in human milk, demonstrate that between 0.01 % and 0.03 % of maternal injected activity of technetium Tc 99m Sestamibi was excreted in human milk. Technetium Tc 99m Sestamibi accumulates in the lactating breast [see *Clinical Considerations*]. There are limited data in the scientific literature on effects of Technetium Tc 99m Sestamibi on the breastfed infant or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Technetium Tc 99m Sestamibi and any potential adverse effects on the breastfed infant from Technetium Tc 99m Sestamibi or from the underlying maternal condition.

Clinical Considerations

Interruption of breastfeeding after exposure of Technetium Tc 99m Sestamibi is not necessary because Technetium Tc 99m Sestamibi excretion in breast milk is low. However, a lactating woman may restrict close contact with her breast fed infant to a maximum of 5 hours in the 24 hour period after Technetium Tc 99m Sestamibi administration in order to minimize radiation exposure.

8.4 Pediatric Use

Safety and effectiveness in the pediatric population have not been established.

No evidence of diagnostic efficacy or clinical utility of Technetium Tc 99m Sestamibi scan was found in clinical studies of children and adolescents with Kawasaki disease.

A prospective study of 445 pediatric patients with Kawasaki disease was designed to determine the predictive value of Technetium Tc 99m Sestamibi rest and stress myocardial perfusion imaging to define a pediatric population with Kawasaki disease that was at risk of developing cardiac events. Cardiac events were defined as cardiac death, MI, hospitalization due to cardiac etiology, heart failure, CABG or coronary angioplasty. The standard of truth was defined as cardiac events occurring 6 months following the administration of Technetium Tc 99m Sestamibi. Only three cardiac events were observed at six months in this study. In all three cases, the scan was negative. No clinically meaningful measurements of sensitivity, specificity or other diagnostic performance parameters could be demonstrated in this study.

A ten year retrospective case history study of pediatric Kawasaki disease patients who completed Technetium Tc 99m Sestamibi myocardial perfusion imaging and who had coronary angiography within three months of the Technetium Tc 99m Sestamibi scan was designed to measure sensitivity and specificity of Technetium Tc 99m Sestamibi scan. Out of 72 patients who had both evaluable Technetium Tc 99m Sestamibi scans and evaluable angiographic images, only one patient had both an abnormal angiogram and an abnormal Technetium Tc 99m Sestamibi scan. No clinically meaningful measurements of sensitivity, specificity or other diagnostic performance parameters could be demonstrated in this study.

In a clinical pharmacology study, 46 pediatric patients with Kawasaki disease received Technetium Tc 99m Sestamibi administration at the following doses: 0.1 to 0.2 mCi/kg for rest, 0.3 mCi/kg for stress in one day studies; 0.2 mCi/kg for rest and 0.2 mCi/kg for stress in two day studies.

The radioactivity both in younger children and in adolescents exhibited PK profiles similar to those previously reported in adults [see *Clinical Pharmacology* (12.3)].

The radiation absorbed doses in adolescents, both at rest and stress, were similar to those observed in adults [see *Dosage and Administration* (2.2)]. When comparing weight-adjusted radioactivity (up to 0.3 mCi/kg) doses administered to adolescents and younger children to the recommended dose administered to adults (up to 30 mCi), the radiation absorbed doses in both adolescents and younger children were similar to those in adults.

Adverse events were evaluated in 609 pediatric patients from the three clinical studies described above. The frequency and the type of the adverse events were similar to the ones observed in the studies of Technetium Tc 99m Sestamibi in adults. Two of the 609 had a serious adverse event: one patient received a Technetium Tc 99m Sestamibi overdose but remained asymptomatic, and one patient had an asthma exacerbation following administration.

8.5 Geriatric Use

Of 3068 patients in clinical studies of Technetium Tc 99m Sestamibi for myocardial imaging, 693 patients were 65 or older and 121 were 75 or older.

Of 673 patients in clinical studies of Technetium Tc 99m Sestamibi for breast imaging, 138 patients were 65 or older and 30 were 75 or older.

Based on the evaluation of the frequency of adverse events and review of vital signs data, no overall differences in safety were observed between these subjects and younger subjects. Although reported clinical experience has not identified differences in response between elderly and younger patients, greater sensitivity of some older individuals cannot be ruled out.

10 OVERDOSAGE

The clinical consequences of overdosing with Technetium Tc 99m Sestamibi are not known.

11 DESCRIPTION

Each 10 mL vial contains a sterile, non-pyrogenic, lyophilized mixture of:

- Tetrakis(2-methoxyisobutylisonitrile)copper(I) Tetrafluoroborate – 1 mg
- Sodium Citrate Dihydrate – 2.6 mg
- L-Cysteine Hydrochloride Monohydrate – 1 mg
- Mannitol – 20 mg
- Stannous Chloride, Dihydrate, minimum (SnCl₂·2H₂O) – 0.025 mg
- Stannous Chloride, Dihydrate (SnCl₂·2H₂O) – 0.075 mg
- Tin Chloride (stannous and stannic) Dihydrate, maximum (as SnCl₂·2H₂O) – 0.086 mg

Prior to lyophilization the pH is 5.6 to 5.8. The contents of the vial are lyophilized and stored under nitrogen.

This drug is administered by intravenous injection for diagnostic use after reconstitution with sterile, non-pyrogenic, oxidant-free Sodium Per technetate Tc 99m Injection. The pH of the reconstituted product is 5.5 (5.0 to 6.0). No bacteriostatic preservative is present.

The precise structure of the technetium complex is Tc 99m[MIBI]₆⁺ where MIBI is 2-methoxyisobutylisonitrile.

11.1 Physical Characteristics

Technetium Tc 99m decays by isomeric transition with a physical half-life of 6.02 hours¹. Photons that are useful for detection and imaging studies are listed below in Table 3.

Table 3. Principal Radiation Emission Data

Radiation	Mean % per Disintegration	Mean Energy (keV)
Gamma-2	89.07	140.5

¹Kocher, David, C., Radioactive Decay Data Tables, DOE/TIC-11026, 108(1981).

11.2 External Radiation

The specific gamma ray constant for Tc 99m is 5.4 microcoulombs/kg-MBq-hr (0.78 R/mCi-hr) at 1 cm. The first half value layer is 0.017 cm of Pb. A range of values for the relative attenuation of the radiation emitted by this radionuclide that results from interposition of various thicknesses of Pb is shown in Table 4. To facilitate control of the radiation exposure from Megabequerel (millicurie) amounts of this radionuclide, the use of a 0.25 cm thickness of Pb will attenuate the radiation emitted by a factor of 1,000.

Table 4. Radiation Attenuation by Lead Shielding

Shield Thickness (Pb) cm	Coefficient of Attenuation
0.017	0.5
0.08	10 ⁻¹
0.16	10 ⁻²
0.25	10 ⁻³
0.33	10 ⁻⁴

To correct for physical decay of this radionuclide, the fractions that remain at selected intervals after the time of calibration are shown in Table 5.

Table 5. Physical Decay Chart; Tc 99m Half Life 6.02 Hours

Hours	Fraction Remaining	Hours	Fraction Remaining
0*	1.000	7	0.447
1	0.891	8	0.398
2	0.794	9	0.355
3	0.708	10	0.316
4	0.631	11	0.282
5	0.562	12	0.251
6	0.501		

*Calibration Time

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Technetium Tc 99m Sestamibi is a cationic Tc 99m complex which has been found to accumulate in viable myocardial tissue in a manner analogous to that of thallous chloride Tl-201. Scintigraphic images obtained in humans after the intravenous administration of the drug have been comparable to those obtained with thallous chloride Tl-201 in normal and abnormal myocardial tissue.

Animal studies have shown that myocardial uptake is not blocked when the sodium pump mechanism is inhibited. Although studies of subcellular fractionation and electron micrographic analysis of heart cell aggregates suggest that Tc 99m Sestamibi cellular retention occurs specifically within the mitochondria as a result of electrostatic interactions, the clinical relevance of these findings has not been determined.

The mechanism of Tc 99m Sestamibi localization in various types of breast tissue (e.g., benign, inflammatory, malignant, fibrous) has not been established.

12.3 Pharmacokinetics

Pulmonary activity is negligible even immediately after injection. Blood clearance studies indicate that the fast clearing component clears with a t½ of 4.3 minutes at rest, and clears with a t½ of 1.6 minutes under exercise conditions. At five minutes post injection about 8 % of the injected dose remains in circulation. There is less than 1 % protein binding of Technetium Tc 99m Sestamibi in plasma. The myocardial biological half-life is approximately six hours after a rest or exercise injection. The biological half-life for the liver is approximately 30 minutes after a rest or exercise injection. The effective half-life of clearance (which includes both the biological half-life and radionuclide decay) for the heart is approximately 3 hours, and for the liver is approximately 30 minutes, after a rest or exercise injection. The ideal imaging time reflects the best compromise between heart count rate and surrounding organ uptake.

Myocardial uptake which is coronary flow dependent is 1.2 % of the injected dose at rest and 1.5 % of the injected dose at exercise. Table 6 illustrates the biological clearance as well as effective clearance (which includes biological clearance and radionuclide decay) of Tc 99m Sestamibi from the heart and liver.

[Organ concentrations expressed as percentage of injected dose; data based on an average of 5 subjects at rest and 5 subjects during exercise].

Table 6. Biological and Effective Clearance

Time	Rest				STRESS			
	Heart		Liver		Heart		Liver	
	Biological	Effective	Biological	Effective	Biological	Effective	Biological	Effective
5 min.	1.2	1.2	19.6	19.4	1.5	1.5	5.9	5.8
30 min.	1.1	1.0	12.2	11.5	1.4	1.3	4.5	4.2
1 hour	1.0	0.9	5.6	5.0	1.4	1.2	2.4	2.1
2 hours	1.0	0.8	2.2	1.7	1.2	1.0	0.9	0.7
4 hours	0.8	0.5	0.7	0.4	1.0	0.6	0.3	0.2

A study in a dog myocardial ischemia model reported that Technetium Tc 99m Sestamibi undergoes myocardial distribution (redistribution), although more slowly and less completely than thallous chloride Tl-201. A study in a dog myocardial infarction model reported that the drug showed no redistribution of any consequence. Definitive human studies to demonstrate possible redistribution have not been reported. In patients with documented myocardial infarction, imaging revealed the infarct up to four hours post dose.

Metabolism

The agent is excreted without any evidence of metabolism.

Elimination

The major pathway for clearance of Tc 99m Sestamibi is the hepatobiliary system. Activity from the gall bladder appears in the intestines within one hour of injection. Twenty-seven percent of the injected dose is excreted in the urine, and approximately thirty-three percent of the injected dose is cleared through the feces in 48 hours.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In comparison with most other diagnostic technetium labeled radiopharmaceuticals, the radiation dose to the ovaries (1.5 rads/30 mCi at rest, 1.2 rads/30 mCi at exercise) is high. Minimal exposure (ALARA) is necessary in women of childbearing capability [see *Dosage and Administration* (2.2)].

The active intermediate, Cu(MIBI)₆BF₄, was evaluated for genotoxic potential in a battery of five tests. No genotoxic activity was observed in the Ames, CHO/HPRT and sister chromatid exchange tests (all *in vitro*). At cytotoxic concentrations (> 20 mcg/mL), an increase in cells with chromosome aberrations was observed in the *in vitro* human lymphocyte assay. Cu(MIBI)₆BF₄ did not show genotoxic effects in the *in vivo* mouse micronucleus test at a dose which caused systemic and bone marrow toxicity (9 mg/kg, > 600 x maximal human dose).

14 CLINICAL STUDIES

CLINICAL TRIALS:

Myocardial Imaging: In a trial of rest and stress Tc 99m Sestamibi imaging, the relationship of normal or abnormal perfusion scans and long term cardiac events was evaluated in 521 patients (511 men, 10 women) with stable chest pain. There were 73.9 % Caucasians, 25.9 % Blacks and 0.2 % Asians. The mean age was 59.6 years (range: 29 to 84 years). All patients had a baseline rest and exercise Tc 99m Sestamibi scan and were followed for 13.2 ± 4.9 months (range: 1 to 24 months). Images were correlated with the occurrence of a cardiac event (cardiac death or non-fatal myocardial infarction). In this trial as summarized in Table 7, 24/521 (4.6 %) had a cardiac event.

Table 7. Cardiac Events

Baseline Scan ^(a)	Proportion of patients with events by scan results ^(a)	Proportion of scan result in patients with events; N=24 ^(a)	Proportion of event-free patients by scan result ^(a)
Normal	1/206 (0.5 %)	1/24 (4.2 %)	205/206 (99.5 %)
Abnormal	23/315 (7.3 %) ^(b)	23/24 (95.8 %) ^(b)	292/315 (92.7 %) ^(b)

(a) Note: Similar findings were found in two studies with patients who had pharmacologic stress Tc 99m Sestamibi imaging.

(b) p < 0.01

Although patients with normal images had a lower cardiac event rate than those with abnormal images, in all patients with abnormal images it was not possible to predict which patient would be likely to have further cardiac events; i.e., such individuals were not distinguishable from other patients with abnormal images.

The findings were not evaluated for defect location, disease duration, specific vessel involvement or intervening management.

In earlier trials, using a template consisting of the anterior wall, inferior-posterior wall and isolated apex, localization in the anterior or inferior-posterior wall in patients with suspected angina or coronary artery disease was shown. Disease localization isolated to the apex has not been established. In adults, Tc 99m Sestamibi has not been studied or evaluated in cardiac disorders other than coronary artery disease.

Breast Imaging: Technetium Tc 99m Sestamibi was evaluated in two multicenter, clinical trials of a total of 673 woman patients. Overall the mean age was 52 (range 23 to 87 years). The racial and ethnic representation was 70 % Caucasian, 15 % African-American, 14 % Hispanic and 1 % Asian. Both clinical studies evaluated women who were referred for further evaluation for either: 1) a mammographically detected (with varying degrees of malignant likelihood) but not palpable breast lesion (study A, n=387, mean age = 54 years), or 2) a palpable breast lesion (study B, n=286, mean age = 50 years). In both studies all patients were scheduled for biopsy.

Technetium Tc 99m Sestamibi (20 to 30 mCi) was injected intravenously in a vein that was contralateral to the breast lesion in question. Planar imaging was completed with a high resolution collimator with a 10 % window centered at 140 keV, and 128 x 128 matrix. An initial marker image, that was not used in the data analysis, was obtained using a cobalt Co57 point source as a marker of a palpable mass. Images were obtained 5 minutes after injection as follows: lateral image of the affected breast for 10 minutes, lateral image of the contralateral breast for 10 minutes, and an anterior image of both breasts for 10 minutes. For the lateral image the patients were positioned in a prone position. For the anterior image, the patients were supine. The Technetium Tc 99m Sestamibi scintigraphic images were read in a randomized method by two groups of three blinded readers. Technetium Tc 99m Sestamibi uptake was scored as: normal (no uptake), equivocal, low, moderate, or high uptake. The results of Technetium Tc 99m Sestamibi images and mammography were analyzed in comparison to histopathologic findings of malignant or non-malignant disease.

As shown in Table 8 for the 483 evaluable patients, the sensitivity and specificity of any degree of Technetium Tc 99m Sestamibi uptake appear to vary with the presence or absence of palpable mass.

Table 8. Overall technetium Tc 99m Sestamibi Blinded Results of Target Lesions^(a) Identified at Study Entry^(b)

STATISTIC	Study A Non-Palpable Mass and an Abnormal Mammogram	Study B Palpable Mass
Number of Patients and Lesions	N=277 Patients with 300 Lesions	N=206 Patients with 240 Lesions
Sensitivity	52 (42,62) ^(c)	76 (67,83)
Specificity	94 (89,96)	85 (77,91)
PPV ^(d)	79 (67,88)	83 (74,89)
NPV ^(d)	80 (74,85)	78 (69,84)
Agreement	80 (75,85)	80 (75,85)
Prevalence	32 (27,37)	49 (43,56)

(a) Excludes all discordant lesions not identified at entry and excludes 25 equivocal interpretations from Study A and 32 equivocal interpretations from Study B (see Tables 9 and 10)

(b) Some patients had more than one target lesion

(c) Median and approximated 95 % Confidence Interval

(d) PPV = Positive Predict Value; NPV = Negative Predict Value

In separate retrospective subset analyses of 259 patients with dense (heterogeneously/extremely dense) and 275 patients with fatty (almost entirely fat/numerous vague densities) breast tissue, the Technetium Tc 99m Sestamibi results were similar. Overall, the studies were not designed to compare the performance of Technetium Tc 99m Sestamibi with the performance of mammography in patients with breast densities or other coexistent breast tissue disorders.

In general the histology seems to correlate with the degree of Technetium Tc 99m Sestamibi uptake. As shown in Tables 9 and 10, the majority of the normal Technetium Tc 99m Sestamibi images are associated with non-malignant tissue (78 % to 81 %) and the majority of low, moderate or high uptake Technetium Tc 99m Sestamibi images are associated with malignant disease (79 % to 83 %). In an individual patient, however, the intensity of Technetium Tc 99m Sestamibi uptake can not be used to confirm the presence or absence of malignancy. Equivocal results do not have a correlation with histology.

Table 9. Degree of Technetium Tc 99m Sestamibi Breast Imaging Uptake in Comparison to Histopathology Results in Patients with Mammographically Detected Non-Palpable Lesions* (Study A)

	Normal Uptake N=249 lesions	Equivocal Uptake N=25 lesions	Low, Moderate or High Uptake N=66 lesions
Non-malignant**	201 (81 %)	14 (56 %)	14 (21 %)
Malignant	48 (19 %)	11 (44 %)	52 (79 %)

* Median finding for 3 blinded readers

** Includes benign tissue, fibroadenoma, benign intramammary nodes, radial scar

Table 10. Degree of Technetium Tc 99m Sestamibi Breast Imaging Uptake in Comparison to Histopathology Results in Patients with Palpable Lesions* (Study B)

	Normal Uptake N=129 lesions	Equivocal Uptake N=32 lesions	Low, Moderate or High Uptake N=115 lesions
Non-malignant**	100 (78 %)	19 (59 %)	20 (17 %)
Malignant	29 (22 %)	13 (41 %)	95 (83 %)

* Median finding for 3 blinded readers

** Includes benign tissue, fibroadenoma, benign intramammary nodes, radial scar

An estimate of the likelihood of malignancy based on the Technetium Tc 99m Sestamibi uptake score in combination with the mammographic score has not been studied.

In these two studies approximately 150 additional, non-biopsied lesions were found to be positive after Technetium Tc 99m Sestamibi imaging. These lesions were identified in sites that did not physically correlate with identified entry criteria mammographic lesions and these lesions were not palpable. These lesions were not biopsied. Whether these lesions were benign or malignant is not known. Technetium Tc 99m Sestamibi uptake can occur in both benign and malignant disease. THE CLINICAL USEFULNESS OF A POSITIVE Technetium Tc 99m Sestamibi IMAGE IN THE ABSENCE OF AN ABNORMAL MAMMOGRAM OR A PALPABLE LESION IS NOT KNOWN.

16 HOW SUPPLIED/STORAGE AND HANDLING

Kit for Preparation of Technetium Tc 99m Sestamibi Injection is supplied as a 10 mL vial in kits of five (5) (NDC # 45548-141-05), ten (10) (NDC # 45548-141-10) and thirty (30) (NDC # 45548-141-30), sterile and non pyrogenic.

The patient dose should be measured by a suitable radioactivity calibration system immediately prior to patient administration. Radiochemical purity should be checked prior to patient administration.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

Prior to lyophilization the pH is between 5.6 and 5.8. The contents of the vial are lyophilized and stored under nitrogen. Store at 15 °C to 25 °C (59 °F to 77 °F) before and after reconstitution. Kit for Preparation of Technetium Tc 99m Sestamibi Injection contains no preservatives. Included in each kit of five (5) vials, ten (10) vials and thirty (30) vials, is a package insert and a sufficient number of vial shield labels and radiation warning labels.

This reagent kit is approved for distribution to persons licensed by the U.S. Nuclear Regulatory Commission to use byproduct material identified in 10 CFR 35.200 or under an equivalent license issued by an Agreement State.

17 PATIENT COUNSELING INFORMATION

Kit for the Preparation of Technetium Tc 99m Sestamibi Injection contains the same active ingredient as CARDIOLITE® and MIRALUMA® which are different names for the same drug (Kit for the Preparation of Technetium Tc 99m Sestamibi Injection). Patients should be advised to inform their health care provider if they had an allergic reaction to either drug or if they had an imaging study with either drug.

Lactation: Interruption of breastfeeding after exposure to Technetium Tc 99m Sestamibi is not necessary, however, a lactating woman should be advised to consider restricting close contact with her breast fed infant to a maximum of 5 hours in the 24 hour period after Technetium Tc 99m Sestamibi administration in order to minimize radiation exposure [see *Use in Specific Populations* (8.2)].

CARDIOLITE®, MIRALUMA® and Recon-o-Stat are trademarks of Lantheus Medical Imaging, Inc.

Manufactured for:
Jubilant DraxImage Inc.,
Kirkland, Quebec, Canada,
H9H 4J4.

For Ordering Tel: Toll Free 888-633-5343

All Other Business: 888-633-5343

Made in Canada